



Abdominal Imaging

D



Mamdouh.m5@gmail.com





This document was generated automatically by Vizle

Your Personal Video Reader Assistant Learn from Videos Faster and Smarter

VIZLE PRO / BIZ

PDF, PPT Watermarks

- Convert entire videos
- Customize to retain all essential content
- Include Spoken Transcripts
- Customer support

Visit https://vizle.offnote.co/pricing to learn more

VIZLE FREE PLAN

PDF only Watermarks


- Convert videos partially
- Slides may be skipped*
- Usage restrictions
- No Customer support

Visit https://vizle.offnote.co to try free

Login to Vizle to unlock more slides*



CT entrography - enteroclysis



- Fluoroscopic placement of a 12–16-French nasojejunal tube
- Contrast material (1500–2500 ml of 0.5% methylcellulose) is infused
- 250 ml per minute.
- An anticholinergic compound (N-butyl hyoscine bromide) is injected
- Non contrast scans are obtained first



This document was generated automatically by Vizle

Your Personal Video Reader Assistant Learn from Videos Faster and Smarter

VIZLE PRO / BIZ

PDF, PPT Watermarks

- Convert entire videos
- Customize to retain all essential content
- Include Spoken Transcripts
- Customer support

Visit https://vizle.offnote.co/pricing to learn more

VIZLE FREE PLAN

PDF only Watermarks

- Convert videos partially
- Slides may be skipped*
- Usage restrictions
- No Customer support

Visit https://vizle.offnote.co to try free

Login to Vizle to unlock more slides*